

# Dynamics, Degradation and Future Challenges of Wetlands in South Sumatra Province, Indonesia

Mustika Edi Armanto<sup>1,\*</sup>, Elisa Wildayana<sup>1</sup>, and Bella Syakina<sup>1</sup>

<sup>1</sup>Faculty of Agriculture, Universitas Sriwijaya, South Sumatra, Indonesia

**Abstract.** This paper aims to analyze dynamics, degradation and future challenges of wetlands in South Sumatra Province, Indonesia. Research methods using Focus Group Discussion (FGD), research data and information are processed using SPSS program. There are four dominant types of wetlands in South Sumatra, namely *lebak* swamps, tidal swamps, rivers, lakes, and peatlands. The area of *lebak* swamps has dominantly increased from year to year, while other wetlands have decreased extensively over the years. All species living in wetlands have decreased in number; the most dominant decline was experienced by fish, gymnosperm and fern and at least in moss and angiosperm. The main causes of wetlands degradation are weak enforcement and legal compliance of inadequate policies, reclamation and excessive drainage. Current efforts to protect the wetlands are restoration, groundwater control, fish population conservation and wetland reserves.

## 1 Introduction

Wetlands with an area of about 3.0 million ha occupy at least 30% of the total area of South Sumatra Province and mostly is located on the East coast of South Sumatra Province [1, 2]. Their uniqueness lies in relation to important twelve elements of the ecosystem, namely sea, beach or coastal, muddy plains, peatlands, *lebak*, lagoon, estuaries, lakes, rivers, islands, coastal forests and mangrove forests. The wetlands are capable of providing minimum 55% of ecosystem services, and have been severely degraded due to excessive reclamation, population pressures and weak enforcement and legal compliance of inadequate policies over the last 40 years. Healthy wetlands ecosystems play an important role in sustainable economic development [3].

Wetlands have been mostly used for oil palm, rubber, fishery, livestock, food agriculture, industry, residential, recreation, shopping centers and others. Changes in the uses of wetlands are mostly due to human intervention [4], policies and concepts of wetland utilization planning, which are often undertaken for economic benefit only [5-7]. Wetland environment aspects as a supporting media for human life are still getting less attention. At the present time the government is trying to restore and rehabilitate; trying to

---

\*Corresponding author: [mediarmanto@unsri.ac.id](mailto:mediarmanto@unsri.ac.id)

minimize wetlands degradation to a minimum level [8]. Most wetlands are owned by individuals, private or local communities who are free to cultivate without regard to the environmental impacts of ecosystem balance. Plantation companies are licensed to cultivate their concession areas. They are free to cut trees including mangroves and damage the natural treasures for economic development activities, such as shrimp and fish ponds [9, 10]. If this situation is continually allowed, it is feared that it will negatively impact the balance of the mangrove ecosystem, peatlands and wetlands as a whole [11].

From a positive point of view, systematic management is able to nurture biodiversity in wetlands, in addition to contributing economically to the local population in particular and society at large. This paper aims to analyze dynamics, degradation and future challenges of wetlands in South Sumatra Province, Indonesia. This paper is expected to be useful as input for the government in making policies for wetlands conservation and restoration. This paper is also capable of providing a proportional form of management in managing wetlands as well as management steps or strategies, constraints and future prospects of wetlands.

## 2 Materials and Methods

The study belonged to the qualitative research conducted in January to May 2018. The research areas are located in the whole South Sumatra Province, Indonesia. Around 100 respondents were taken by purposive sampling and interviewed by using semi-structured and face-to-face approach, consisting of open questions to enable perception developing.



**Fig. 1.** Research location in South Sumatra Province (Source: Google)

Primarily research data were directly collected from respondents of local communities by using methods of Focus Group Discussion (FGD), in-depth study, field observation, direct interview. Data collecting from beneficiaries and stakeholders was done through governmental expert meetings; classrooms where students consist of public officials; projects obtained from private companies; activities of road shows; presentation of government and private seminars etc. Respondents from scientists and researchers were interviewed through presentations of national and international seminars; media appearance; recent appointments from key stakeholders. Secondary data were taken from research and research reports, and reviewing documents related to this research.

## 3 Results and Discussions

Wetlands are spread unevenly among the region of South Sumatra; namely (1) the South-Eastern region is dominated by *lebak* swamps (fresh water swamps) and peatlands; (2) various types of lakes, rivers and *lebak* swamps are found in the Western part; (3) the North East regions are dominated by tidal swamps, and rivers; (4) the Eastern region is dominated by tidal wetlands, rivers, peatlands; (5) Coastal areas are covered by tidal swamps, river estuaries, and swamp mud; (6) the middle region is covered by *lebak*

swamps, river systems and lakes; (7) the lower parts of middle region is performed by the lake and river system; and (8) the most western region is occupied by lakes and groups of highland mountain (Figure 1).

### 3.1 Dynamics of Wetlands in South Sumatra

There are five types of dominant wetlands located in the South Sumatra Province region, namely *lebak* swamps, tides, rivers, lakes and peatlands. These five lands have undergone many changes in land area over the last 40 years. This changes occurs due to natural factors (i.e. forest and land fires, floods, earthquake, geologic tectonic) or human intervention (land conversion, economic development activities, and others. The wetland losses or expand over the last 40 years in South Sumatra is presented in Table 1.

**Table 1.** The wetland losses or expand over the last 40 years in South Sumatra.

Wetland type	Acreages of area (%)		Area losses or expand (%)
	1990	2017	
Lebak swamps	30.55	33.43	+9.42
Tidal swamps	20.89	22.29	+6.70
Rivers	11.56	12.34	+6.74
Lakes	12.78	13.98	+9.38
Peatlands	13.45	10.32	-23.27
Unidentified	10.77	7.64	-29.06
Total	100.00	100.00	-20.09

Source: Field survey results (2018), Landsat images (1992; 2017)

There are four types of wetlands that have increased and the order of increasing area in percentage over the last 40 years are as follows: *lebak*; lakes; rivers and tidal swamps with the following values in sequence are as follows 9.42%, 9.38%, 6.74%, and 6.70% respectively. An area increase is induced by natural and human intervention. During the rainy season, other wetlands such as peatlands are flooded, so it becomes *lebak* and tidal swamps. This causes *lebak* and tidal percentage area has increased for the last 40 years.

Decreasing peatlands will be in the percentage of -23.27% due to natural factors and human intervention. Peatlands often experience fires and floods. Fires occur during the dry season where the soil becomes dry, so does the vegetation on the peatlands. This situation triggers easy fire occurs. Fires on peatlands can also occur due to unintentional, but fatal consequences, where people often carelessly do land clearing by burning. On dry land and hot weather, the little flame coming from land clearing is getting bigger and surely spreading and widespread.

Another natural factor affecting the decline of peatlands is flood. Floods generally occur during the rainy season where the water drops in a large volume and lasts longer or even daily. This resulted in floods on peatlands. Peatlands are experiencing a very high water inundation to cause peatlands is not visible. In addition to fires and floods, the area of peatlands is also reduced due to the land conversion process. Many peatlands are undergoing land conversion, the average peatlands is transformed into plantation land. There are several plantation companies that use peatlands into plantations.

Unidentified wetlands also decreased by around -29.06% due to human intervention factor. In 1990 humans did not yet have the capabilities and tools that could be used to identify wetlands. So the percentage of unidentified wetlands area at that time reached 10.77%. However, development of the human era begins to learn and have the tools to identify wetlands. Currently humans have used many tools to identify wetlands like Drone.

This tool is a tool developed by humans that can be used to see the state of land and land area effectively and efficiently.

An increase in the percentage of lake area around 9.38% is caused by human intervention factor. Humans do a lot of developments improving the welfare for the region. The lake is one of the natural beauties that much preferred by the community and as one of the center of family tourism. This condition becomes a potential to increase lake area for making the lake as a tourist attraction.

*Lebak* swamps belong to the largest freshwater wetlands and are increasingly widespread from year to year (around 9.42%). Based on hydro topography, height and length of inundation, *lebak* swamps are divided into five typologies, namely shallow; *pematang*; middle; deep; and *lebaklebung* (Table 2). Management based on *lebak* typology alone cannot give optimal results. This typology approach is useful for agronomic purposes only, but it cannot be used for integrated land management because of the high and long standing factors varies depending on the season and the dynamics of hydrology. So it is not surprising that if *lebak* reclamation is very intensive, then *lebak* typology rapidly turns into other *lebak* typologies or wide composition of each typology quickly changed (Table 2).

**Table 2.** Lebak swamp typology and alternative uses

Typology, their positions & area	Hydrological characters	Landuse in seasons	
		Rain	Dry
Shallow (located along embankment of rivers, alluvial plains and flat topography); 25.23%	Very dynamic, flooded (<3 months/year) in rainy season, inundation height <25 cm, and dry in dry season)	Rice, plantation, livestock, forestry, industry, housing	Fallow, forestry, plantation, livestock, industry, housing
<i>Pematang</i> (on embankment of rivers, river valley and flat topography); 18.78%	Dynamic, flooded <3 months/year, inundation height 25-50 cm, and dry in the dry season	Rice, plantation, livestock, forestry	Rice, forestry plantation, livestock,
Middle (on flood plains and river valleys); 22.02%	Dynamic, flooded 3-6 months/year, and inundation height 50-100 cm	Fishery	Rice, fallow
Deep (located next to floodplain and basin); 21.79%	Less dynamic, flooded 6-9 months/year, inundation height 100-150 cm, and peat <50 cm	Fishery, fallow	Fishery, fallow
<i>LebakLebung</i> (in deep basin); 12.18%	Static, always inundated, inundation height > 150 cm, and peat 50-100 cm	Fishery, fallow	Fishery, fallow

Source: Field survey results (2018), Landsat images (1992; 2017)

Shallow *lebak* is very dynamic means that any *lebak* development will directly affect the extent of shallow *lebak* because *lebak* development has a direct impact on the increase of sedimentation which resulted in the widespread area of the flooded river basin. Because the nature is very dynamic, then shallow *lebak* area can be smallest or widest. Another case with *lebaklebung* is static compared to middle *lebak* due to permanent and deep inundation.

### 3.2 Biodiversity Loss

Wetlands are under threat from reclamation, excessive drainage and irrigation, dam, building, pollution, excessive use of resources, biological invasion and global climate change. Their impacts are biodiversity loss, land and water pollution, overfishing, and

sedimentation. This threat should be minimized to preserve the wetlands. Wetlands occupy only around 30% of the total area of South Sumatra Province, but are able to provide a large number of ecosystem services, including fresh water supplies, flood arrangements, wastewater purification, natural resources, wildlife habitats, and marine life preservation (Table 3).

**Table 3.** The predicted numbers of species living in wetlands of South Sumatra

Taxon	Numbers of species		Losses (%)
	1990	2017	
Natural Vegetation			
Gymnosperm	268	197	-26.49
Fern	2.367	2.128	-10.11
Moss	2.249	2.124	-5.56
Angiosperm	31.000	28.077	-9.43
Subtotal	35.884	32.526	-9.37
Animals			
Fish	3.017	2.191	-27.38
Bird	1.256	1.077	-14.25
Mammal	532	428	-12.50
Reptilian	318	249	-21.70
Amphibian	271	214	-21.03
Subtotal	5.394	4.159	-22.90

Sources: Compiled from BPS (2015); Field survey results (2018); Google (2017); Forestry Department (2015) and other literature

**Gymnosperm.** This population reduction is due to illegal logging by irresponsible parties. Gymnosperm especially acacia and pine trees are widely used for furniture, pulp and paper and common industry. Because of these, it is encouraging certain parties to do illegal logging without thinking about environmental sustainability. **Fern** population decline is due to people living around wetlands utilizing ferns for daily consumption of food and medicine. Since ferns commonly grow and live wildly in bushes or forest, people take ferns for free, thus why its population is decreasing.

**Moss** population reduction is due to the global warming effect. The growth of moss tends to be slower in dry and infertile areas because moss belongs to epiphytic vegetation that is less suitable to live in hot and dry places. The effect of global warming inhibits moss growth, so that the population is decreasing. **Angiosperm** is the largest group of vegetation and very important for human life because it is useful as a source of food for humans and animals. Some examples of angiosperm are fruits and flowers (such as mango, apple, tomato, banana, durian, orange, orchid, sunflower, rice and others). Angiosperm is widely used as a source of human and animal food, land conversion to settlement and industry.

**Fish** populations decreased due to overfishing using poisonous and electrical illegal instruments. Global warming also affects the increase in temperature and increased acidity levels due to increased carbon dioxide. This has an impact on the decline of fish populations, because warming temperatures in the water will inhibit the growth of fish and even at risk of fish death. Throwing garbage carelessly into the waters is causing water pollution that threatens the life of fish inside. **Amphibians** decline populations is caused by the low availability of animal food resources in the form of small fish. Less source of food makes amphibians cannot meet their food needs affecting their inhibition of growth and development. The reduction in amphibian populations is also due to habitat destruction and climate change.

**Birds** reduction population is due to the growing human activity damaging the bird habitats, for example land clearing of forest for settlements and industry where the forest is a natural habitat of birds, humiliation of birds and depletion of bird food stocks. In accordance with the order of the food chain if the amphibians or small fish are extinct, then the bird as a consumer will also go extinct because in general birds must eat at least half of their weight every day. **Reptiles** presence is very important in natural food networks as predators and prey. As reptile predators will prey on other animals such as fish, frogs and birds to support their survival. Reduced reptile food sources will certainly increase the risk of declining reptile populations. Reptile extinctions are also triggered by climate change threatening their existence as well as habitat destruction and excessive extraction for animal trade. **Mammals** belong to the generalist species eating many types of food are far more able to survive in a changing environment.

In addition, it increases the frequency of floods from once every five years or ten years. The loss of wetlands also resulted in the extinction of many species requiring wetlands as their vital habitat. Among known animal species that have been extinct in the last 50 years, most of them are wetlands species, including large head fish (*Aspiorhynchuslaticeps*), estuarine crocodiles (*Crocodylusporosus*), betook fish, tapa, *tengkeleso*, elephant and wild deer (*Elaphurusdavidianus*). Most extinction is caused by habitat loss due to wetlands damage. The disappearance of wild rice (*Oryza meyeriana*), which was found in the South Sumatra swamps of South Sumatra in the 1980s and used to create hybrid rice; These new crops increase rice production, but wild rice populations disappear from their natural environment. In addition to the direct destruction of wetlands, there are other issues related to the health and conservation of wetlands, such as water pollution, loss of C storage, and biological invasion. This problem is as bad as the loss of wetlands.

### 3.3 Determinant Degradation Causes of Wetlands

Many factors contributed to degradation of wetlands, including high population pressure on natural resources, lack of understanding of wetlands values, weak enforcement and legal compliance of inadequate policies, weak regulation and environmental regulations, excessive drainage and irrigation and rapid economic growth. Here, we discuss three major causes of wetlands loss over the years: reclamation, misguided policy and midlanduses, and excessive drainage and irrigation.

#### 3.3.1 Reclamation

Reclamation is a major cause of loss of wetlands. Most reclamation has been going on for the last 40 years. Around 133.500 ha of wetlands have been used for plantations, agriculture, fish ponds, settlement and industry obtained from wetlands conversion. Overall, reclamation alone could reach 78% of total wetlands losses. If landscape of wetlands is evaluated, wetlands are expanding around + 32.24%, while wetlands loss is about -52.33%, thus the total wetlands lost as much as -20.09% for 27 years (1990-2017) or about 0.74% per year due to erosion and soil compaction (Table 1). Most eroded soils are transported into general waters (rivers, sea, lakes, basins, valley and so on).

#### 3.3.2 Misguided Policy and Mislanduses

The recovery capacity loss of wetlands is so high. It was primarily the result of reclamation policies supported by the government during the last 40 years. Efforts for food resilience is the main driving force of government reclamation policy, and large-scale reclamation by

building embankments and water canals networks requires greater financial and labor support. Only the government can provide it. However, recent natural disasters bring the attention of governments and society to the severe environmental problems caused by the policy of economic growth by all means. The realization of the environmental consequences of misguided policies also leads to changes in attitudes toward the environment in general and wetlands in particular. As a result, large-scale reclamation is still permitted by the government, although wetlands and natural swamps are under threat from environmental impacts, as many local governments continue to regard them as potential wetlands resources for exploitation.

### **3.3.3 Excessive Drainage and Irrigation**

Decreased groundwater levels are a major cause of loss of natural wetlands. Drainage development for plantation, agricultural and industrial development has led to a reduction in water flow to wetlands over the last 40 years. The drained wetlands area increased by around 340% from 1990 to 2017, and most of the water was lost due to excessive drainage (120-150%) and low irrigation efficiency (15-35%). The excessive use of water by agriculture and other industries resulted in a low river flow at the bottom of the river having no water flow. Beside that around 75% of water dams and water gates do not function properly, thus they are not able to manage water according to plant needs and also disrupt the species migration route waters and finally alter the water fauna.

## **3.4 Current Efforts to Protect Wetlands**

Wetlands are protected under a three-class system: wetlands reserves (full protection from development and human activities), wetlands parks (full protection from development, but used for ecotourism), and beautiful gardens (protection from development, but open to the public for recreation). Some important actions should be taken by the government in protecting, restoring, and creating wetlands:

### **3.4.1 Peatland Restoration**

South Sumatra began to restore the degraded wetlands in early 2014s by establishing Peat Restoration Agency (BRG). BRG functions are to protect and restore existing wetlands, to recover lost wetlands, and to tackle other wetlands problems. BRG do also how to restore the water quality of natural peatlands, and to restore pollution purification capacity in wetlands. The new policy has been made to restore some peatlands of reclaimed farms to wetlands or native peatlands, namely groundwater management; fish population conservation; and wetland reserves and restorations.

Groundwater management plans are implemented to improve water control in the fields and to protect wetlands, particularly to treat highly degraded wetlands. The action plan focuses on the control of point and nonpoint sources of problems, water management and water control by involving local society. In addition, many laws and regulations on water management has been enacted by national, provincial and local governments.

Regulations to ban or prohibit fishing in the reproductive period have been carried out, for example at the mouth of the river and fishing grounds. Action plans for restocking (replenishing) wetlands with fish seedlings from hatcheries have begun, including both commercial and endangered species of fish, have been released into common waters (lakes, rivers, tidal, and peatlands). Wetlands reserves consist of core areas, where human activity is prohibited to cultivate agriculture, and the areas are determined as buffering zones, where

some human activities are only permitted by permission. The government began to build wetlands natural reserves in early 2000s.

## 4 Conclusions

The results and discussion of this research can be drawn some conclusions as follows: Dominant wetlands in South Sumatra are namely *lebak*, tidal, rivers, lakes, peatlands, and unidentified. *Lebak* swamps belong to dominant wetlands changing dominantly each year followed by tidal swamps, rivers, and lakes while other types decrease continuously. Predicted species living in wetlands are decreasing in number; the most dominant decrease was experienced by fish, gymnosperm, mammal, reptilian, bird, amphibian, fern, angiosperm and moss respectively. The major causes of wetlands loss over the years are reclamation, misguided policy and mislanduses, and excessive drainage and irrigation. Current efforts to protect wetlands are peatland restoration, groundwater management, fish population and conservation and wetland reserves and restorations.

## Acknowledgments

The authors would like to acknowledge the financial assistance provided by Universitas Sriwijaya along with the highly valued staff inputs at the Faculty of Agriculture of Universitas Sriwijaya. We would like to thank all students for their comments on early versions of this paper along with valuable feedback from anonymous reviewers.

## References

1. Sarno, R.A. Suwignyo, Z. Dahlan, Munandar, M.R. Ridho, N. Aminasih, Harmida, M.E. Armanto, E. Wildayana. *Biodiversitas* **18**, 3 (2017)
2. M.E Armanto,, M.A. Adzemi, E. Wildayana, M.S. Imanudin. *J. of Sustainability Science and Management* **8**, 1 (2013)
3. M.E Armanto,. E. Wildayana. *J. of Wetlands Environmental Managements* **4**, 1 (2016)
4. S.M. Lu. *J. of Sustainable Development* **103** (2017)
5. K. Fobissie, D. Etongo, M. Kanninen. *Journal of Sustainable Development* **10**, 5 (2017)
6. E. Wildayana, M.E. Armanto. *J. of Sustainable Development* **11**, 3 ( 2018)
7. E. Wildayana, M.E. Armanto. *J. of Sustainable Development* **11**, 2 (2018)
8. M.E Armanto., M.S. Imanudin, E. Wildayana, H. Junediand M. Zuhdi. *Sriwijaya Journal of Environment* **1**, 3 (2016)
9. E. Wildayana. *J. of Agricultural Science* **23**, 6 (2017)
10. E. Wildayana, M.E. Armanto. *Bulgarian Journal of Agricultural Science.* **24**, 2 (2018)
11. I. Zahri, D. Adriani, E. Wildayana, Sabaruddin, M.U. Harun, *Bulgarian Journal of Agricultural Science.* **24**, 2 (2018)