

Analysis of exposure of inhabitants of Polish cities to air pollution with particulate matters with application of statistical and geostatistical tools

Izabela Sówka^{1,*}, Alicja Nych¹, Dominik Kobus², Yarolsav Bezyk¹, and Maciej Zathej³

¹Wrocław University of Science and Technology, Faculty of Environmental Engineering, Wyb. Wyspińskiego 27, 50-370 Wrocław, Poland

²inFAIR, www.infair.eu, dominik.kobus@infair.eu

³Wrocław University of Science and Technology, Faculty of Architecture, ul. Bolesława Prusa 53/55, 50-317 Wrocław, Poland

Abstract. The aim of this research was the analysis of air quality in selected Polish cities in 2008-2018 combined with health assessment based on the methodology and software of World Health Organization (AirQ +) as well as an analysis of main directions of activities aimed at improving air quality in Polish cities. The results of calculations and spatial analysis of exposure indicate the year 2010 as the one with the highest concentrations of PM₁₀ and PM_{2.5}, as well as Małopolskie and Śląskie Voivodships (agglomerations: Upper Silesia, Krakow and Rybnik-Jastrzębska) as the two most threatened by negative impact on human life and health. In a detailed analysis of the Wrocław agglomeration, for 2017, the estimated number of deaths of adults caused by short-term exposure to PM_{2.5} with a limiting concentration above 10 µg/m³ (in accordance with WHO guidelines on air quality) was over 130 people, and the estimated number of patients hospitalized because of cardiovascular and respiratory diseases due to the above exposure was 160 and 80 cases, respectively. The analysis showed more than 80 cases of asthma attacks in children with asthma caused by short-term exposure and three death cases in infants due to long-term exposure to PM₁₀.

1 Introduction

Air pollution is one of the most important determinants of health. Numerous epidemiological studies have demonstrated the relationship between air pollution and a wide range of adverse health effects in general population; effects ranged from subtle subclinical effects to premature death. Some groups – for example, elderly people, children, pregnant women and people with chronic illnesses (e.g. asthma) may be more at risk and may be more at risk from exposure to air pollution. In addition, some groups may be exposed to a higher level of outdoor air pollution, for example people living near busy traffic routes or in specific occupations or socio-economic groups [1].

* Corresponding author: izabela.sowka@pwr.edu.pl

In 2010, lung diseases were the cause of 6.9% of all hospitalizations in Poland, and death rates in 2010 due to respiratory illnesses amounted to 50.6 per 100,000 residents [2]. Because of general cardiovascular diseases, on the other hand, over one million people were treated [3]. Also, in the document "Policy paper for health care for 2014–2020" [4] published in 2014, respiratory illnesses were listed among the most important health problems of Polish citizens. They were also included in the group of health issues that generated the largest incapacity expenses in 2010 (7.5% of the total expenditure) [5]. According to the 2012 State Health Report [6], mortality due to cardio-pulmonary syndrome and other heart diseases (51.5 thousand deaths in 2010 – 30% of deaths due to the whole circulatory causes), is higher in Poland than throughout the European Union. Death rates in 2010 due to cardiorespiratory syndrome and other heart problems were in total 134.9 per 100,000. residents [2].

The aim of this study was to analyse air quality in selected Polish cities, combined with a health assessment based on the methodology and software of World Health Organization (AirQ+).

2 Research methodology

2.1 Air pollution monitoring data analysis

The analysis of variability of the concentrations of selected air pollutants (PM₁₀ and PM_{2.5}) was carried out for urban agglomerations with populations of over 250,000, according to the classification adopted in the Regulation of the Minister of Environment of August 2, 2012 on zones in which air quality is assessed (Journal of Laws from 2012, item 914). Results from measurement stations included in the National Environmental Monitoring network, recorded in the period between 2008 and 2018 were taken into account and Air Quality Portal of the Chief Inspectorate for Environmental Protection was the source of data used for calculations. Data from 2018 should be treated as not finally approved yet by bodies responsible for monitoring.

The analysis took into account data from an urban and suburban background and industrial stations (located within agglomeration boundaries). Due to the significantly limited spatial representativeness, uneven distribution in agglomerations and lack of continuity in the period covered by the analysis, data from traffic stations was omitted. Prior to the processing of data, they were selected due to the quality level, as well as applied measurement methodologies. Annual series of measurement results with a completeness of at least 75%, obtained by the manual gravimetric method (being the reference method), as well as automatic methods in the absence of manual measurements in a given location were included. For the purposes of analysis, the 1-hour data series were aggregated to 24-hour data, with the condition of minimum 75% completeness of data in the day. Then, for each agglomeration, daily indicators for individual pollutants were calculated: daily average and maximum daily mean values for each day for all stations in the agglomeration. For each agglomeration an annual index was calculated, being an arithmetic mean of daily average indicators.

2.2 WHO model

The analysis of the impact (both short- and long-term) of selected air pollutants on health and lives of exposed people was carried out using AirQ+ software developed by WHO. The effect of exposure to excessive occurrence in atmospheric air of particulate matter (PM_{2.5} and PM₁₀) has been analysed. In particular, for 12 agglomerations throughout Poland, the effect of long-term PM_{2.5} exposure on the total mortality was analysed. In addition, for the same agglomerations, the effect of exposure to PM_{2.5} on cases of hospital admission for

cardiovascular (HACD) and respiratory diseases (HARD) was evaluated. Results of concentrations of individual pollutants from the period between 2008 and 2018 were used for calculations. Additionally, for the Wrocław agglomeration, for 2017, analysis of long-term impact from PM₁₀ dust on post neonatal infant mortality and short-term impact on the occurrence of asthma attacks in children with asthma was carried out. Moreover, effects of short-term PM_{2.5} exposure on adults mortality, as well as HACD and HARD were analysed. Statistical data on population, deaths and baselines for hospitalization in the analysed areas was taken from the Central Statistical Office and Ministry of Health. The default values in the software were used for relative risk (which estimates the relationship between exposure and health problems and indicates the probability of disease development in the exposed group, compared to those who are not exposed), as well as the value of the limit concentration below which health impact is not analysed. Table 1 presents the relative risk for each health effect analysed in this paper.

Table 1. Relative risk (95% confidence interval) and used for investigating PM health effects.

Health effect	Relative risk (95% CI)
Short term effect PM _{2.5} Mortality, all causes (adults age 30+ years)	1.0123 (1.0045–1.0201)
Short term effect PM _{2.5} Hospital admission for cardiovascular diseases	1.0091 (1.0017–1.0166)
Short term effect PM _{2.5} Hospital admission for respiratory diseases	1.008 (1.0048–1.0112)
Short term effect PM ₁₀ Incidence of asthma symptoms in asthmatic children	1.028 (1.006–1.051)
Long term effect PM ₁₀ Post neonatal infant Mortality, all cause	1.04 (1.02–1.07)
Long term effect PM _{2.5} Mortality, all causes (adults age 30+ years)	1.062 (1.04–1.083)

2.3 Geostatistical analysis

A geostatistical analysis is a very useful approach that allows providing accessible information related to the ambient air quality monitoring network. Then, the geographical information system (GIS) as a set of powerful techniques of geoprocessing, mapping and spatial analysis is capable in terms of distribution and spatial patterns for each air pollutants [7].

The GIS estimated air pollution surfaces can be certainly useful for further analysis of the air pollution series in the urban area at different time scales and prediction purposes of air pollutants for exposure studies. GIS was used in the work to visualize the results of statistical analyses of estimated number of attributable cases of total mortality caused by long-term exposure on PM_{2.5} in 2010 against the average of 2008–2018 and the estimated number of attributable cases of patients hospitalized for cardiovascular (HACD) and respiratory (HARD) diseases in Poland in 2010 caused by short-term exposure on PM_{2.5}.

3 Research results and discussion

3.1 Characteristics of dust pollution in selected Polish cities

Cities in Poland, including the largest agglomerations, show very high diversity in terms of air quality. This also applies to particulate matter, the concentration of which was analysed. The areas with lower concentrations of this pollution include the agglomerations: Trójmiejska and Szczecińska, while the highest concentrations occur in Rybnicko-Jastrzębska, Górnośląska and Krakowska. The annual variability course of PM₁₀ and PM_{2.5} concentrations indicates a declining trend, with the occurrence of peaks in 2010 in most cities (Fig. 1. – example for PM_{2.5}). This year was characterized, among other things, by such meteorological conditions in winter, that are conducive to an increased emission of

particulate matter, mainly from the domestic heating sources – e.g. a large number of days with the minimum temperature below 10°C [8]. Such conditions, combined with low wind speed and the occurrence of the temperature inversion phenomenon, lead to an increase in emission and weakening dispersion conditions and as a consequence to higher pollution. The occurrence of smog episodes is quite common in autumn and winter in Poland, including in the analysed agglomerations [9–11].

In the last four years, a stable situation with respect to the average concentration of PM_{2.5} can be observed in most cities, even with increases in some agglomerations.




Fig. 1. Variability of annual PM_{2.5} indices in agglomerations in 2008–2018 (source: own elaboration based on data from the State Environmental Monitoring).

PM₁₀ and PM_{2.5} concentrations for all cities show large seasonal, weekly and daily variations. An example of data from 2017 for one of the stations located in the Wrocław Agglomeration is shown in Figure 2. January and February were characterized by very high concentrations, including deep episodes – average concentrations significantly differed in this period from the rest of the year. The concentrations on weekend days are noticeably lower, which can be attributed to smaller car traffic that impacts this station. Such an influence is also indicated by analysis of daily variation, where traffic peaks are clearly visible, but they are clearer (especially in the morning) for the larger fraction of measured dust (PM₁₀).




Fig. 2. Seasonal, weekly and daily PM₁₀ concentration variability at the Wrocław-Korzeniowskiego station in 2017 (source: own state-owned monitoring based on the State Environmental Monitoring).




Fig. 4. Estimated number of attributable cases of patients hospitalized for cardiovascular (HACD) and respiratory (HARD) diseases in Poland in 2010 caused by short-term exposure on $PM_{2.5}$.

3.4 Analysis of main directions of activities aimed at improving air quality in Polish cities

Formal activities related to the improvement of air quality in Polish cities depend mainly on legal solutions and subsequent financing of their effects. In this context, cities use the law laid down by the Polish national parliament and regional councils. Local activities, being a commune or urban initiative, are complementary. Actions such as: a day without the car, tree planting or dissemination of information about threats resulting from exceeding limit values of air pollution are primarily aimed at educating and shaping social attitudes. However, they do not eliminate the causes of pollution.

In order to achieve the observance of environmental quality standards, in Poland, based on the Act – Environmental Protection Law (Journal of Laws 2001 No. 62 item 627 with later amendments), air protection programs adopted by way of a resolution by voivodship parliaments are created. They contain recommendations being an obligation addressed mainly to local governments and, therefore, also to cities. Unfortunately, this mechanism is ineffective as evidenced by the state of atmospheric pollution and the need to introduce additional restrictions. Therefore, on the basis of the amendment to the Act – Environmental Protection Law, which was passed on November 12, 2015, regional councils gained the possibility of introducing restrictions and prohibitions on the operation of installations in which fuels are burned - adoption of the so-called anti-smog resolutions (Table 3).

In parallel to the preparation of the anti-smog resolutions, social pressure has caused the preparation of system solutions at a central level in Poland. The ordinance of the Minister of Development and Finance on requirements for solid fuel boilers has been adopted, which formally eliminated from 01.07.2018 the sale of boilers that do not comply with EN 303-5: 2012. The second significant regulation, issued by the Minister of Energy, defines the quality requirements for solid fuels. It uses a two-year grace period for the use of coal fines below 3 mm in diameter. In this respect, the national provision is more liberal than regional solutions. This results in maintaining a trade in fuels whose combustion is prohibited by regulations adopted by the regional councils. This paradox sustains illegal combustion of high-emission solid fuels by 30 June 2020.

Table 3. A list of voivodships in which the regional councils have introduced restrictions and prohibitions regarding the operation of installations in which fuels are burned pursuant to art. 96 of the Act – Environmental Protection Law (drawn up on the basis of resolutions of regional councils included in the analysis).

	Voivodship	Separate regulations for cities	The date of the most stringent prohibition	Requirements for fuels and installations
1.	dolnośląskie		Resolution adopted on 30.11.2017	
		Wrocław	01.07.2028	The target total elimination of solid fuels.
		Spas	01.07.2028	Target total elimination of solid fuels in seven locations, in four locations permitted installations in accordance with EC Regulation (EU) 2015/1185 with regard to ecodesign requirements.
2.	lubuskie		Resolution adopted on 18.06.2018	
		Gorzów Wielkopolski	01.01.2023	Class 5 installations according to PN-EN 303-5: 2012.
		Zielona Góra	01.01.2023	Class 5 installations according to PN-EN 303-5: 2012.
3.	łódzkie		Resolution adopted on 24.10.2017	
4.	małopolskie		Resolution adopted on 23.01.2017	
		Kraków	01.09.2019	The target total elimination of solid fuels.
5.	mazowieckie		Resolution adopted on 24.10.2017	
6.	opolskie		Resolution adopted on 26.09.2017	
7.	podkarpackie		Resolution adopted on 23.04.2018	
8.	śląskie		Resolution adopted on 07.04.2017	
9.	wielkopolskie		Resolution adopted on 18.12.2017	
		Poznań	01.01.2028	Installations in accordance with the EC Regulation (EU) 2015/1185 with regard to ecodesign requirements.
		Kalisz	01.01.2028	Class 5 installations according to PN-EN 303-5: 2012
10.	zachodniopomorskie		Resolution adopted on 26.09.2018	

4 Conclusions

The analysis of research results and calculations carried out indicate higher concentrations of dust contamination, which may contribute to the deterioration of the quality of life of the inhabitants of Polish cities. Reduction of air pollution in Polish cities should therefore become a priority in activities in the field of environmental protection in Poland.

A real improvement of air quality in cities will be possible only after the transformation of heating systems and a widespread use of zero-emission or low-emission heat production technology with a simultaneous improvement of energy efficiency in housing, housing estates and transport systems. For this purpose, a radical change in the energy policy of the state is needed. In this context, the energy policy and the air quality policy pursued by regions and cities will be a secondary activity. Programs for co-financing changes in heat sources, which are mainly responsible for air pollution, should be considered as the initial stage of converting the heat protection system in Poland.

The investigations were co-financed within the framework of the order No. 0401/00581/18.

References

1. WHO Regional Office for Europe, *Health risk assessment of air pollution – general principles* (Copenhagen, 2016)
2. Ministerstwo Zdrowia, *Program Polityki Zdrowotnej* (2017)
3. Ministerstwo Zdrowia, *Narodowy Program Wyrównywania Dostępności do Profilaktyki i Leczenia Chorób Układu Sercowo-Naczyniowego POLKARD na lata 2013–2016* (2014)
4. Ministerstwo Zdrowia, *Krajowe ramy strategiczne* (2015)
5. E. Karczewicz, A. Kania, *Wydatki na świadczenia z ubezpieczeń społecznych związane z niezdolnością do pracy w 2010 r.* (2012)
6. B. Wojtyński, P. Goryński, B. Moskalewicz, *Sytuacja zdrowotna ludności Polski i jej uwarunkowania* (pod red. NIZP-PZH, Warszawa, 2011)
7. J. Gulliver, K. de Hoogh, D. Fecht, D. Vienneau, D. Briggs, *Atmos. Environ.*, **45**, 7072–7080 (2011)
8. A. Chlebowska-Styś, I. Sówka, D. Kobus, Ł. Pachurka, *E3S Web of Conferences* **17**, 1–8 (2017)
9. D. Kobus, A. Nych, I. Sówka, *E3S Web of Conferences* **44**, 00070 (2018)
10. J. Iwanek, J. Strużewska, J. Kamiński, P. Durka, D. Kobus, J. Kostrzewa, T. Pecka, *Analiza wybranych epizodów wysokich stężeń pyłu PM10 z lat 2013–2016, Etap II, Epizody z lat 2015–2016* The work done for CIEP (Instytut Ochrony Środowiska – Państwowy Instytut Badawczy: IOŚ PIB, 2017)
11. K. Juda-Rezler, B. Toczko (ed.), *Pyły drobne w atmosferze. Kompendium wiedzy o zanieczyszczeniu powietrza pyłem zawieszonym w Polsce* (Inspekcja Ochrony Środowiska: IOŚ, Biblioteka Monitoringu Środowiska, Warszawa, 2016)
12. Rozporządzenie Ministra Rozwoju i Finansów z dnia 1 sierpnia 2017 r. w sprawie wymagań dla kotłów na paliwo stałe
13. Rozporządzenie Ministra Energii z dnia 27 września 2018 r. w sprawie wymagań jakościowych dla paliw stałych
14. Ustawa – Prawo ochrony środowiska Dz. U. 2001 Nr 62 poz. 627 z późn. zm.